

Breaking of $B - L$ in superstring inspired E_6 model.

A. Kartavtsev

Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany

E-mail: akartavt@het.physik.uni-dortmund.de

Abstract. In the framework of the superstring inspired E_6 model, ‘low-energy’ extensions of the standard model compatible with leptogenesis are considered and masses of right-handed neutrinos in two scenarios allowed by long-lived protons are discussed. The presence of two additional generations allows breaking of $B - L$ without generating nonzero vacuum expectation values of right-handed sneutrinos of the three known generations. After the symmetry breaking, right-handed neutrinos acquire Majorana masses $M_{\nu^c} \simeq 10^{11}$ GeV. Within the framework of a simple discrete symmetry, assumptions made to provide a large mass of right-handed neutrinos are shown to be self-consistent. Supersymmetric structure of the theory ensures that large corrections, associated with the presence of a (super)heavy gauge field, cancel out.

1. Introduction

The observed baryon asymmetry of the universe is one of the most interesting problems of particle physics and cosmology. As was suggested by Sakharov, the cosmological baryon excess can be generated through the baryon number violating decays of heavy particles [1]. Any microscopic theory able to reproduce the observed asymmetry must satisfy Sakharov's three conditions:

- baryon (or baryon minus lepton) number non-conservation;
- C and CP violation;
- deviation from thermal equilibrium.

The condition of baryon number violation makes a strong appeal to physics beyond the standard model (SM). It has been proven to be difficult to generate an excess of baryons through direct violation of the baryon number. A very attractive scenario of generating lepton number asymmetry through lepton number violating decays of heavy right-handed neutrinos, which is later converted into the baryon number by sphalerons [2, 3], has been suggested by Fukugita and Yanagida [4]. This scenario requires any viable extension of the SM to contain heavy right-handed neutrinos, which, furthermore, are the reason for small masses of conventional neutrinos.

Augmented with three right-handed neutrinos, the SM itself fails to reproduce the observed baryon asymmetry of the universe Y_B due to the relatively small CP violation and too weak phase transition. Electroweak baryogenesis in the minimal supersymmetric standard model (MSSM) is not yet ruled out, but is close to being so [5]. Since a significant amount of asymmetry is generated during the electroweak (EW) phase transition, from the practical point of view it is of interest to consider simple 'low-energy' extensions of the SM (i.e., models with additional $U(1)$ or $SU(2)$ gauge symmetry and extended particle content) able to reproduce the observed Y_B [6, 7, 8]. From the theoretical point of view, it is interesting to investigate lepto- and baryogenesis in grand unification theories (GUT). In this paper, an attempt to derive 'low-energy' extensions of the SM, compatible with the Fukugita–Yanagida scenario, from a supersymmetric GUT model is made.

The choice of a supersymmetric model is motivated by two reasons. First, scalar superpartners of right-handed neutrinos can be used to break down $B - L$ symmetry spontaneously by the Higgs mechanism. Second, supersymmetry (if it is broken only softly, as is usually assumed) provides cancellation of large quantum corrections associated with the presence of high scales in the theory.

A minimal gauge group with right-handed neutrino contained in the same representation as the 15 known fermions is $SO(10)$. An additional representation is required to fit Higgs fields responsible for breaking of the electroweak symmetry. Following the ideas of grand unification, it seems more natural to choose a gauge group which at the unification scale treats all states on an equal footing, i.e., which contains the known states along with right-handed neutrino and Higgs doublets in the fundamental

representation. This is the case for the exceptional group E_6 , one of the most prominent candidates for a unified theory. Its characteristic features are as follows.

- It naturally follows from breaking of superstring $E'_8 \otimes E_8$ with $N_f \mathbf{27} + \delta(\mathbf{27} + \overline{\mathbf{27}})$ chiral superfields and $\mathbf{78}$ vector superfields as zero mode spectra.
- It allows chiral representations and its fundamental representation fits the fifteen known fermions along with right-handed neutrino and two Higgs doublets.
- Gauge anomalies are automatically canceled.

In section 2, the particle content of the fundamental representation and possible charge assignments are discussed. There are six charge assignments, compatible with the SM. Charge conservation in the processes involving states of different generations requires that the same charge assignment be used for all generations.

Some of the fields in the adjoint representation may lead to a rapid proton decay. Constraints implied by a long-lived proton are considered in section 3. In addition to the intermediate gauge groups listed in [9] the $SU(5) \times U^2(1)$ is allowed for two charge assignments. Nevertheless, Yukawa interactions implied by the residual $SU(5)$ symmetry make the rapid proton decay mediated by new down-type quarks unavoidable, unless those are very heavy.

Section 4 discusses $B - L$ symmetry breaking and masses of the right-handed neutrinos. An interesting feature of the model under consideration is the presence of additional $\delta(\mathbf{27} + \overline{\mathbf{27}})$ generations, scalar right-handed neutrinos of which may be used to break $B - L$ symmetry spontaneously. Introduced simple discrete symmetry ensures that $B - L$ is broken at the scale which is sufficiently high for generating large masses of the right-handed neutrinos and that right-handed scalar neutrinos of the three known generations do not acquire a vacuum expectation value (VEV). It also forbids Yukawa couplings which, if present, would induce large masses of the conventional neutrinos. The supersymmetric structure of the theory ensures that large quantum corrections to masses of scalars, associated with the presence of heavy gauge fields, cancel out. After the $B - L$ breaking, the ‘low-energy’ gauge group is $SU_C(3) \times SU_L(2) \times U^2(1)$.

Finally, section 5 is devoted to some concluding remarks.

2. Particle content and charge assignments

The ten-dimensional $E_8 \times E'_8$ heterotic superstring theory compactifies to the $M_4 \times \Gamma$ and yields a low-energy theory with $N = 1$ supersymmetry. Γ is the Calabi–Yau manifold with $SU(3)$ holonomy. If Γ is simply connected, then $E_8 \times E'_8$ gauge group breaks down to the $E_6 \times E'_8$ subgroup. For a multiply connected manifold the initial gauge group breaks down to $G \times E'_8$ where G is a subgroup E_6 [10]. In this scheme, chiral superfields $N_f \mathbf{27} + \delta(\mathbf{27} + \overline{\mathbf{27}})$ and $\mathbf{78}$ vector superfields of E_6 emerge as the zero mode spectra. For a wide class of models $\delta = 1$. States in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ are denoted by χ and $\overline{\chi}$ respectively, whereas states in $N_f \mathbf{27}$ are denoted by ψ .

The fundamental representation of E_6 fits the 15 known states of the SM along with two Higgs doublets (H^u and H^d), a pair of down-type quarks (D and D^c) and two SM singlets ν^c and S .

Table 1. Particle content of E_6 : fundamental representation.

The Standard Model states					New states					
$B-L$	Y	I_3	Q_{em}	P	$B-L$	Y	I_3	Q_{em}	P	
1/3	1/3	1/2	2/3	u	} Q	0	1	1/2	1	E^c
1/3	1/3	-1/2	-1/3	d		0	1	-1/2	0	N^c
-1	-1	1/2	0	ν	} L	0	-1	1/2	0	N
-1	-1	-1/2	-1	e		0	-1	-1/2	-1	E
-1/3	-4/3	0	-2/3	u^c	-2/3	-2/3	0	-1/3	D	
-1/3	2/3	0	1/3	d^c	2/3	2/3	0	1/3	D^c	
1	2	0	1	e^c	1	0	0	0	ν^c	
					0	0	0	0	S	

A systematic study of quantum numbers of states in the fundamental and adjoint representations is conveniently performed using the Cartan–Weyl method [11, 12]. In the framework of this method, each state is represented by its weight. Weights of **27** and **78** of E_6 are presented in tables 7 and 8, respectively. The hypercharge of any state is given by a scalar product of the hypercharge operator Y and the weight of the state. One can easily check that the hypercharge assignment is not unique — there are three assignments which reproduce the SM:

$$3Y_1 = (1, \bar{1}, 1, \bar{3}, \bar{1}, 0), \quad (1a)$$

$$3Y_2 = (1, \bar{1}, \bar{5}, \bar{3}, \bar{1}, 0), \quad (1b)$$

$$3Y_3 = (1, \bar{1}, 1, 3, \bar{1}, 0). \quad (1c)$$

There are also three $B - L$ operators, i.e. three assignments, which reproduce $B - L$ quantum numbers of the SM states:

$$3B - L_1 = (1, \bar{1}, \bar{2}, \bar{3}, \bar{1}, 0), \quad (2a)$$

$$3B - L_2 = (1, \bar{1}, 1, 0, \bar{1}, 0), \quad (2b)$$

$$3B - L_3 = (1, \bar{1}, \bar{2}, 0, \bar{1}, 0). \quad (2c)$$

Different assignments correspond to different embeddings of states into subgroups of E_6 . Out of nine $Y_i \times B - L_j$ combinations six are compatible with the SM:

$$\text{I : } (Y_1, B - L_1), \quad \text{II : } (Y_1, B - L_2), \quad \text{III : } (Y_2, B - L_1)$$

$$\text{IV : } (Y_2, B - L_3), \quad \text{V : } (Y_3, B - L_2), \quad \text{VI : } (Y_3, B - L_3)$$

Since $B - L$ is gauged, the Majorana mass of the right-handed neutrino, which is an essential ingredient of leptogenesis, is forbidden unless $B - L$ symmetry is broken down. Present data on neutrino masses as well as theoretical estimates of leptogenesis in other GUT models favour the $10^{10} - 10^{12}$ GeV ν^c mass range [8, 13].

Using weights, one can derive the form of trilinear interactions following the algorithm developed in [14]. In the flavor basis

$$\begin{aligned}
 W = & \lambda_1^{ijk} u_i^c(Q_j H_k^u) + \lambda_2^{ijk} d_i^c(Q_j H_k^d) + \lambda_3^{ijk} e_i^c(L_j H_k^d) + \lambda_4^{ijk} S_i(H_j^u H_k^d) + \lambda_5^{ijk} S_i D_j D_k^c \\
 & + [\lambda_6^{ijk} e_i^c u_j^c D_k + \lambda_7^{ijk} D_i^c(Q_j L_k) + \lambda_8^{ijk} d_i^c \nu_j^c D_k] + [\lambda_9^{ijk} D_i(Q_j Q_k) + \lambda_{10}^{ijk} D_i^c u_j^c d_k^c] \\
 & + \lambda_{11}^{ijk} \nu_i^c(L_j H_k^u)
 \end{aligned} \quad (3)$$

Using table 7, one can check that the corresponding weights add up to zero at each vertex. The form of the superpotential is independent of the charge assignment in the sense that the change of assignment will only result in ‘permutation’ of vertices. See assignments I, II and III, for example:

	$[10\bar{1}001]$	$[00010\bar{1}]$	$[\bar{1}01\bar{1}00]$	$[1\bar{1}01\bar{1}0]$	$[01\bar{1}010]$	$[101\bar{1}00]$
I	ν^c	ν	N^c	S	N	N^c
II	S	N	N^c	ν^c	ν	N^c
III	e^c	ν	E	S	E^c	E

Along with vertices with $i = j = k$ (i.e. states of the same generation) superpotential (3) necessarily contains terms with the states of different generations in one vertex. If the charge assignment for one of the generations differs from that for other generations, then this results in nonzero sum of charges in a vertex. For instance, if in the example above assignment II is used for the first weight and assignment I is used for the second and the third weights then

$$\begin{aligned}
 \nu^c \rightarrow S & \Rightarrow \nu^c (\nu N^c) \rightarrow S (\nu N^c), \quad \sum B - L = -1 \\
 S \rightarrow \nu^c & \Rightarrow S (NN^c) \rightarrow \nu^c (NN^c), \quad \sum B - L = 1
 \end{aligned}$$

If instead of assignment II, assignment III is used for the first weight, then it is a sum of electric charges, which is nonzero. This suggests that the same charge assignment should be used for all generations.

3. Gauge mediated proton decay.

Since $SU(5)$ is a subgroup of E_6 , the gauge sector of the model contains X and Y bosons, which are known to mediate proton decay. There are also new gauge fields leading to a rapid proton decay (table 2).

Table 2. Gauge fields which mediate proton decay.

$B - L$	Y	I_3	Q_{em}	P	$B - L$	Y	I_3	Q_{em}	P
2/3	5/3	1/2	4/3	$\bar{X}$	-2/3	1/3	1/2	2/3	$u_{\bar{2}}$
2/3	5/3	-1/2	1/3	Y	-2/3	1/3	-1/2	-1/3	$d_{\bar{2}}$

To assure that the proton is long-lived, these fields must be very heavy — of order of 10^{15} GeV or more. If the only source of masses of those particles is the VEV of neutral

scalars, then the masses of (u_2, d_2) gauge bosons are determined by $\langle \tilde{\nu}^c \rangle$, Y boson mass is of the order of EW symmetry breaking scale and the X bosons remain massless even after all neutral scalars develop a nonzero VEV.

Consequently, the gauge superfields which mediate proton decay have to become massive at the first stage of symmetry breaking. If the manifold Γ is multiply connected, then the effective Higgs mechanism [15, 16] breaks symmetry at the compactification scale and induces large masses of the gauge fields.

From the discussion above it follows that after $E_8 \times E'_8$ breaking the gauge group of the model is not E_6 itself, but a subgroup G of E_6 . A very elaborate analysis of many possible breaking chains has been performed in [9, 16]. It was argued there that a gauge field gets mass of the order of $O(10^{18})$ GeV if $(Z, \rho) \neq 0$. Here Z is the zero root breaking $SU_C(3) \times SU_L(2) \times U_Y(1)$ preserving direction, and ρ is the weight of the gauge field. Scalar products (Z, ρ) for fundamental and adjoint representations of E_6 are listed, respectively, in tables 7 and 8.

Table 3. (Z, ρ) products for $(\mathbf{3}, \mathbf{2})_{\bar{5}}$ gauge fields.

SU(5)	$SU(3) \times SU(2) \times U(1)$	I	II	III	IV	V	VI	(Z, ρ)
24	$(\mathbf{3}, \mathbf{2})_{\bar{5}}$	X	X	d_2	d	d_2	d	α
24	$(\mathbf{3}, \mathbf{2})_{\bar{5}}$	Y	Y	u_2	u	u_2	u	α

The requirement of $(Z, \rho) \neq 0$ for the gauge fields which mediate proton decay does not allow $G = SO(10) \times U(1)$. $(\mathbf{3}, \mathbf{2})_{\bar{5}}$ states (and their charge conjugate) are the only states in the adjoint of $SU(5)$ which are not automatically massless. As seen from table 3, the requirement that these fields be massless (i.e. that $SU(5)$ is unbroken) does not lead to a rapid gauge mediated proton decay for the charge assignments IV and VI. In other words, in addition to the intermediate gauge groups listed in [9] as allowed, $G = SU(5) \times U^2(1)$ is allowed for the charge assignments IV and VI. Nevertheless, in this case the residual $SU(5)$ symmetry implies that couplings in (3) are related in the following way;

$$\lambda_1 = \lambda_2 = \lambda_4 = \lambda_{10}, \quad \lambda_5 = \lambda_7 = \lambda_9, \quad \lambda_3 = \lambda_6, \quad \lambda_8 = \lambda_{11}$$

and the rapid proton decay mediated by heavy down-type quarks D and D^c is unavoidable.

Apart from scenarios with an extended color group there are only two options left:

$$G_{U(1)} = SU_C(3) \times SU_L(2) \times U(1) \times U(1) \times U(1) \quad (4)$$

$$G_{SU_R(2)} = SU_C(3) \times SU_L(2) \times SU_R(2) \times U(1) \times U(1) \quad (5)$$

If the intermediate scale symmetry group is given by (4), then the gauge sector of the SM is supplemented by two neutral states (the rest are superheavy), denoted by ϕ^0 and ω^0 (see table 8).

If the intermediate scale symmetry group is given by (5), then the gauge sector of the SM is supplemented by one $SU_R(2)$ singlet ω^0 and one $SU_R(2)$ triplet $(\phi^c, \phi^0, \bar{\phi}^c)$.

There are three possible $SU_R(3) \rightarrow SU_R(2)_Y$ projections which result in one left–right symmetric (R) and two skew left–right symmetric (R' and R'') models. $SU_R(2)$ counterparts of ϕ^0 for different choices of the projection and the charge assignment are given in table 4.

Table 4. $SU_R(2)$ counterparts of ϕ^0 . Isospin and hypercharge (as well as $B - L$, unless otherwise indicated) of the states designated like the SM states are the same as charges of their SM counterparts. If present, subscript is trice the $B - L$ charge.

	Charge assignment					
	I	II	III	IV	V	VI
R	$e_0^c, \bar{e}_0^c$	$e^c, \bar{e}^c$	$e_0^c, \bar{e}_0^c$	$e^c, \bar{e}^c$	$\nu^c, \bar{\nu}^c$	$\nu^c, \bar{\nu}^c$
R'	$e^c, \bar{e}^c$	$e_0^c, \bar{e}_0^c$	$\nu^c, \bar{\nu}^c$	$\nu^c, \bar{\nu}^c$	$e_0^c, \bar{e}_0^c$	$e^c, \bar{e}^c$
R''	$\nu^c, \bar{\nu}^c$	$\nu^c, \bar{\nu}^c$	$e^c, \bar{e}^c$	$e_0^c, \bar{e}_0^c$	$e^c, \bar{e}^c$	$e_0^c, \bar{e}_0^c$

While $SU_L(2)$ in (5) coincide with that of the standard model, U_{Y_L} is not the SM $U_Y(1)$. Hypercharge Y , as well as $B - L$, is a linear combination of Q_{Y_L} , Q_{Y_R} and I_{3R} . An explicit form of gauge interactions $g_\alpha \Lambda_\alpha [\psi^\dagger T^\alpha \psi]$ for the first charge assignment and left–right symmetric model can be read off from table 7:

$$\begin{aligned}
& \frac{g_{Y_L}}{2} Y_L \left[\bar{Q}Q - 2\bar{D}D - (\bar{H}^u H^u + \bar{H}^d H^d) - \bar{L}L + 2(\bar{e}^c e^c + \bar{\nu}^c \nu^c) + 2\bar{S}S \right] \\
& + \frac{g_{Y_R}}{2} Y_R \left[-(\bar{d}^c d^c + \bar{u}^c u^c) + 2\bar{D}^c D^c + (\bar{H}^u H^u + \bar{H}^d H^d) \right. \\
& \left. - 2\bar{L}L + (\bar{e}^c e^c + \bar{\nu}^c \nu^c) - 2\bar{S}S \right] + \frac{g_{W_R}}{2} W_R^i \left[(d^c, u^c)^\dagger \tau_i (d^c, u^c) \right. \\
& \left. + (H^u, H^d)^\dagger \tau_i (H^u, H^d) + (e^c, \nu^c)^\dagger \tau_i (e^c, \nu^c) \right] + \dots
\end{aligned} \tag{6}$$

At the unification scale the relation among the gauge couplings is as follows:

$$g_{W_R} = \sqrt{3} g_{Y_L} = \sqrt{3} g_{Y_R} = g_{E_6} \tag{7}$$

Instead of Y_L , Y_R and W_R^0 one can use their linear combination. For instance, in the limit (7) linear combination $Y_{B-L} = \frac{1}{\sqrt{2}}(Y_L + Y_R)$ is a gauge field of $U_{B-L}(1)$. For the later purposes it is useful to choose the linear combinations ω'_0 , ϕ'_0 , γ'_0 in such a way, that only one of the new fields interacts with right–handed neutrinos and one of the fields interacts with neither of the SM singlets. After the Standard Model singlets develop VEVs, ω'_0 and ϕ'_0 acquire masses, while γ'_0 , which corresponds to $U_Y(1)$, remains massless. Rewritten in terms of these fields, (6) takes the form

$$\begin{aligned}
& g_{\gamma'_0} \gamma'_0 \frac{1}{6} \left[\bar{Q}Q - 4\bar{u}^c u^c + 2\bar{d}^c d^c + 2\bar{D}^c D^c - 2\bar{D}D + 3\bar{H}^u H^u - 3\bar{H}^d H^d - 3\bar{L}L + 6\bar{e}^c e^c \right] \\
& + g_{\phi'_0} \phi'_0 \left[-\bar{Q}Q - \bar{u}^c u^c - 2\bar{d}^c d^c + 3\bar{D}^c D^c + 2\bar{D}D + 2\bar{H}^u H^u + 3\bar{H}^d H^d \right. \\
& \left. - 2\bar{L}L - \bar{e}^c e^c - 5\bar{S}S \right] + g_{\omega'_0} \omega'_0 \left[\bar{Q}Q + \bar{u}^c u^c - 2\bar{d}^c d^c + \bar{D}^c D^c - 2\bar{D}D \right. \\
& \left. - 2\bar{H}^u H^u + \bar{H}^d H^d - 2\bar{L}L + \bar{e}^c e^c + 4\bar{\nu}^c \nu^c + \bar{S}S \right] + \dots
\end{aligned} \tag{8}$$

At the unification scale gauge couplings $g_{\gamma'_0}$, $g_{\phi'_0}$, $g_{\omega'_0}$ are given by

$$g_{\gamma'_0} = \sqrt{\frac{3}{5}} g_{E_6}, \quad g_{\phi'_0} = \frac{g_{E_6}}{\sqrt{40}}, \quad g_{\omega'_0} = \frac{g_{E_6}}{\sqrt{24}} \quad (9)$$

Reexpressed in terms of ω'_0 , ϕ'_0 , γ'_0 gauge interactions of neutral fields are obviously given by (8) irrespectively of the intermediate scale symmetry group G , charge assignment or particular choice of $SU_R(3) \rightarrow SU_R(2)$, Y projection.

4. Large Majorana masses of the right-handed neutrinos.

There are two SM singlets whose scalar superpartners may be used to break symmetry down to the SM: S and the right-handed neutrino.

The former has zero $B - L$ charge, while the later has $B - L = 1$. Therefore, it is the VEV of the scalar superpartner of the right-handed neutrino that breaks $B - L$ symmetry. S couples to Higgs doublets and its VEV $\langle \tilde{S} \rangle$ is the origin of the μ -term: $\mu = \lambda_4 \langle \tilde{S} \rangle$.

If the right-handed sneutrino which develops the VEV couples to states of the three known generations, it induces huge Dirac masses for components of L and H^u doublets via the last term in (3). Neglecting the possibility that one of the ν^c superfields is decoupled from the other states of the three known generations, one comes to the conclusion that all $\langle \tilde{\nu}^c \rangle = 0$, and the $B - L$ symmetry is broken spontaneously by nonzero $\langle \tilde{\chi}_{\nu^c} \rangle$ and $\langle \tilde{\bar{\chi}}_{\nu^c} \rangle$ and, consequently, that χ_{ν^c} and $\bar{\chi}_{\nu^c}$ are zero modes.

According to [16] chiral superfields in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ can be massive through the Yukawa coupling $\mathbf{27} \cdot \overline{\mathbf{27}} \cdot \mathbf{78}$. If $(Z, \rho) \neq 0$ for a component of $\mathbf{27}$ or $\overline{\mathbf{27}}$ with weight ρ , the corresponding chiral superfield gets compactification scale mass, while N_f $\mathbf{27}$ chiral superfields remain massless.

Table 7 shows, that for both discussed intermediate gauge groups $G_{U(1)}$ and $G_{SU_R(2)}$ and any charge assignment, it is possible to have massless χ_{ν^c} and $\bar{\chi}_{\nu^c}$. In the case of $G_{U(1)}$ right-handed neutrinos χ_{ν^c} and $\bar{\chi}_{\nu^c}$ are the only massless states in $\delta(\mathbf{27} + \overline{\mathbf{27}})$.

In the case of $G_{SU_R(2)}$ the number of zero modes in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ depends on the charge assignment and the particular choice of $SU_R(3) \rightarrow SU_R(2)$ projection (table 5).

Table 5. States in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ which remain massless after the compactification. States of $\mathbf{27}$ and $\overline{\mathbf{27}}$ are labeled here by the same symbol. For instance, ν^c stands for both χ_{ν^c} and $\bar{\chi}_{\nu^c}$. Components of $SU_R(2)$ doublets are put into brackets.

Charge assignment						
I	II	III	IV	V	VI	
R	(ν^c, e^c)	$\nu^c, (H^u, L)$	(ν^c, e^c)	$\nu^c, (H^u, L)$	(ν^c, S)	(ν^c, S)
R'	$\nu^c, (H^u, L)$	(ν^c, e^c)	(ν^c, S)	(ν^c, S)	(ν^c, e^c)	$\nu^c, (H^u, L)$
R''	(ν^c, S)	(ν^c, S)	$\nu^c, (H^u, L)$	(ν^c, e^c)	$\nu^c, (H^u, L)$	(ν^c, e^c)

If supersymmetry is exact, there are no negative mass squared terms needed to break $U_{B-L}(1)$ down spontaneously by Higgs mechanism.

$$m_\chi^2 |\tilde{\chi}_{\nu^c}|^2 + m_{\tilde{\chi}}^2 |\tilde{\tilde{\chi}}_{\nu^c}|^2 - m_{\nu^c ij}^2 \tilde{\nu}_i^{c*} \tilde{\nu}_j^c, \quad |m_\chi^2|, |m_{\tilde{\chi}}^2|, |m_{\nu^c ij}^2| \sim m_{soft}^2 \quad (10)$$

These terms are assumed to come from the E_8' sector, where supersymmetry is considered to break down spontaneously. In the gravity-mediated supersymmetry breaking scenario the magnitude of the soft terms in the visible sector should be roughly of the order of $m_{soft} \sim \langle F \rangle / M_{Pl}$. For the commonly accepted value $m_{soft} \sim 10^3$ GeV the scale of supersymmetry breaking in the hidden sector $\langle F \rangle^{\frac{1}{2}}$ is about 10^{11} GeV. It is interesting to note that this value is of the same order as the desired mass scale of the right-handed neutrinos. One can consider this as a hint that at the stage when right-handed neutrinos acquire masses, the temperature is still high enough to produce them thermally.

$SU_C(3) \times SU_L(2) \times U^3(1)$ model. As is well known, the scalar potential consists of an F -term and D -term coming from the chiral superfield trilinear couplings and the gauge interactions respectively. The renormalizable superpotential (3) does not contain terms relevant for the symmetry breaking. The scalar potential coming from gauge interactions (8) and soft supersymmetry breaking is of the form

$$V = \frac{g_{\omega'_0}^2}{2} \left[\tilde{\psi}^* T_{\omega'_0} \tilde{\psi} + q_{\nu^c} \tilde{\chi}_{\nu^c}^* \tilde{\chi}_{\nu^c} - q_{\nu^c} \tilde{\tilde{\chi}}_{\nu^c}^* \tilde{\tilde{\chi}}_{\nu^c} \right]^2 - [m_\chi^2 |\tilde{\chi}_{\nu^c}|^2 + m_{\tilde{\chi}}^2 |\tilde{\tilde{\chi}}_{\nu^c}|^2] + m_{\nu^c ij}^2 \tilde{\nu}_i^{c*} \tilde{\nu}_j^c \quad (11)$$

On the one hand the VEVs of right-handed sneutrinos of $\delta(\mathbf{27} + \overline{\mathbf{27}})$ are expected to be at least of the same order as masses of ν^c , i.e. $\langle \tilde{\chi}_{\nu^c} \rangle, \langle \tilde{\tilde{\chi}}_{\nu^c} \rangle \geq 10^{11}$ GeV; on the other hand, such a huge VEV should not generate large masses of scalar superpartners via the first term in (11). Consequently, symmetry breaking should occur in the D -flat direction $\langle \tilde{\chi}_{\nu^c} \rangle = \langle \tilde{\tilde{\chi}}_{\nu^c} \rangle$.

Combined with the requirement that all $\langle \tilde{\nu}^c \rangle$ be zero, this means that contribution of the first term in (11) vanishes. To have symmetry breaking by the Higgs mechanism in this direction, the sum of the mass parameters in the second term should be positive: $m_\chi^2 + m_{\tilde{\chi}}^2 > 0$

Non-renormalizable terms arise due to the interactions with exchange of superheavy fields, which correspond to excitations of internal degrees of freedom [18]. The general form of the non-renormalizable superpotential is

$$\begin{aligned} W = & M_c^{-1} [a_1^{ij} \nu_i^c \nu_j^c \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} + a_2^i \nu_i^c \chi_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} + a_3 \chi_{\nu^c} \chi_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c}] \\ & + M_c^{-3} [b_1 \chi_{\nu^c} \chi_{\nu^c} \chi_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} + b_2^i \nu_i^c \chi_{\nu^c} \chi_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} \\ & + b_3^{ij} \nu_i^c \nu_j^c \chi_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} + b_4^{ijk} \nu_i^c \nu_j^c \nu_k^c \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c} \bar{\chi}_{\nu^c}] + \dots \end{aligned} \quad (12)$$

Given that all $\langle \tilde{\nu}^c \rangle$ are zero, only $M_c^{3-2n} (\chi_{\nu^c} \bar{\chi}_{\nu^c})^n$ terms in (12) are relevant for the analysis of symmetry breaking. These terms are invariant with respect to $\chi_{\nu^c} \leftrightarrow \bar{\chi}_{\nu^c}$ transformation, whereas soft supersymmetry breaking terms in (11) are not, unless $m_\chi^2 = m_{\tilde{\chi}}^2$. If this condition is not satisfied, then $\langle \tilde{\chi}_{\nu^c} \rangle$ can not be equal to $\langle \tilde{\tilde{\chi}}_{\nu^c} \rangle$.

Nevertheless, since gauge coupling g is of the order of unity, while m_{soft}/M_c is of many orders of magnitude smaller than unity, the deviation from the D -flat direction $\langle \tilde{\chi}_{\nu^c} \rangle = \langle \tilde{\bar{\chi}}_{\nu^c} \rangle$ is very small. Considering $\langle \tilde{\bar{\chi}}_{\nu^c} \rangle - \langle \tilde{\chi}_{\nu^c} \rangle$ as a small perturbation in (15) one finds

$$\langle \tilde{\bar{\chi}}_{\nu^c} \rangle - \langle \tilde{\chi}_{\nu^c} \rangle \simeq \frac{m_{\bar{\chi}}^2 - m_{\chi}^2}{8g^2 v_0} \quad (13)$$

This effect can be entirely neglected in the analysis of non-renormalizable superpotential (12). As for gauge interactions, such a deviation from the D -flat direction will result in the generation of masses of scalars $\sim m_{\bar{\chi}}^2 - m_{\chi}^2$ via the first term in (11) which add to mass terms coming from the soft supersymmetry breaking. To simplify the analysis $m_{\bar{\chi}}^2$ and m_{χ}^2 are taken to be equal in what follows.

If the first non-vanishing terms in (12) are $M_c^{-1}(\nu^c \bar{\chi}_{\nu^c})^2 + M_c^{3-2n}(\chi_{\nu^c} \bar{\chi}_{\nu^c})^n$ then generated VEV and masses of right-handed neutrinos ν^c are

$$\langle \tilde{\chi}_{\nu^c} \rangle = \langle \tilde{\bar{\chi}}_{\nu^c} \rangle \sim (m_{soft} M_c^{2n-3})^{\frac{1}{2n-2}}, \quad M_{\nu^c} \sim (m_{soft} M_c^{n-2})^{\frac{1}{n-1}} \quad (14)$$

so that M_{ν^c} is of the order of 10^3 GeV for $n = 2$ and is of the order of 10^{11} GeV for $n = 3$.

Large masses of right-handed neutrinos suggest that $n = 3$ and, consequently, $a_3 = 0$ in (12). Moreover, a_2^i and b_2^i , as well as coefficient c_2^i of the (not indicated) next similar term in (12), are zero to avoid nonzero $\langle \tilde{\nu}^c \rangle$ and Dirac-type mass for ν^c . The discrete symmetry of the compactified manifold possibly accomplishes these conditions [19] as well as the absence of bare mass terms $M_{\chi_{\nu^c}} \bar{\chi}_{\nu^c}$ and $M_{\nu^c} \bar{\chi}_{\nu^c}$ in the superpotential (3). There is no reason to expect scale M to be below compactification scale M_c so that the presence of the first term would make spontaneous breaking of $B - L$ by χ_{ν^c} and $\bar{\chi}_{\nu^c}$ impossible, while the presence of the second makes ν^c a component of the super heavy Dirac neutrino. Just as there are no $M_{\chi_{\nu^c}} \bar{\chi}_{\nu^c}$ and $M_{\nu^c} \bar{\chi}_{\nu^c}$ terms, there are no soft supersymmetry breaking terms $b_{\chi\bar{\chi}} \tilde{\chi}_{\nu^c} \tilde{\bar{\chi}}_{\nu^c}$ and $b_{\nu^c\bar{\chi}} \tilde{\nu}^c \tilde{\bar{\chi}}_{\nu^c}$ in (11). If present, the contributions of b_3 and b_4 terms are small and neglected in the following discussion. Then the classical potential is of the form

$$\begin{aligned} V = & 4M_c^{-2} u^2 \varrho_i \varrho_j [a_{li} a_{lj}^* u^2 + a_{ij} a_{nm}^* \varrho_n \varrho_m] + 9b_1 b_1^* M_c^{-6} u^4 v^4 [u^2 + v^2] \\ & + 6M_c^{-4} [b_1^* a_{ij} + b_1 a_{ij}^*] \varrho_i \varrho_j v^3 u^3 + \frac{g^2 q_{\nu^c}^2}{2} [\varrho_i \varrho_i + v^2 - u^2]^2 \\ & - [m_{\chi}^2 v^2 + m_{\bar{\chi}}^2 u^2] + m_{\nu^c}^2 \varrho_{ij} \varrho_i \varrho_j \end{aligned} \quad (15)$$

with $\varrho_i = \langle \tilde{\nu}_i^c \rangle$, $v = \langle \tilde{\chi}_{\nu^c} \rangle$, $u = \langle \tilde{\bar{\chi}}_{\nu^c} \rangle$. For zero ϱ_i the D -flat direction is defined by $u^2 = v^2$ and the minimum of the potential (15) corresponds to

$$v_0 = \sqrt[8]{M_c^6 (m_{\chi}^2 + m_{\bar{\chi}}^2) / (90|b_1|^2)}, \quad M_{\nu^c}^{ij} = 2a_1^{ij} v_0^2 M_c^{-1} \quad (16)$$

For nonzero ϱ_i D -flat direction is defined by $u^2 = v^2 + \varrho_i \varrho_i$. The set of products $\varrho_i \varrho_j$ being considered as parameters, the minimizing of (15) with respect to v gives v as function of $\varrho_i \varrho_j$. True vacuum corresponds to the set $\varrho_i \varrho_j$ which minimizes $V(v(\varrho_i \varrho_j), \varrho_i \varrho_j)$.

Expanding in powers of $\varrho_i \varrho_j$ in the vicinity of $\varrho_i \varrho_j = 0$ and having in mind that partial derivative with respect to v is zero:

$$V(v(\varrho_i \varrho_j), \varrho_i \varrho_j) \simeq V(v_0, 0) + (m_{\nu^c}^2 - m_{\tilde{\chi}}^2) \varrho_i \varrho_j \quad (17)$$

$$+ v_0^4 M_c^{-2} \left[4a_1^{li} a_1^{*lj} + 6(b_1^* a_1^{ij} + b_1 a_1^{*ij}) v_0^2 M_c^{-2} + 45b_1 b_1^* v_0^4 M_c^{-4} \delta^{ij} \right] \varrho_i \varrho_j$$

Since v_0/M_c ratio is small while v_0^2/M_c is large, the derivative (17) is dominated by the first term in square brackets which is a positive definite. Therefore, $\varrho_i = 0$ is at least a local minimum of the potential (15) for a wide range of parameters.

For large $\varrho_i \varrho_j$ it is sufficient to keep only the first term in (15), and an explicit calculation shows that $V(v(\varrho_i \varrho_j), \varrho_i \varrho_j)$ grows with growing $\varrho_i \varrho_j$, i.e. $\varrho_i \varrho_j = 0$ is a global minimum of the classical potential:

$$V(v(\varrho_i \varrho_j), \varrho_i \varrho_j) = -M_c^2 (m_{\tilde{\chi}_{\nu^c}}^2 + m_{\chi_{\nu^c}}^2)^2 (16a_1^{li} a_1^{*lj} \varrho_i \varrho_j)^{-1} + (m_{\tilde{\chi}_{\nu^c}}^2 \delta^{ij} + m_{\nu^c}^2) \varrho_i \varrho_j \quad (18)$$

A discrete symmetry which allows nonzero a_1^{ij} , b_1 and forbids nonzero a_2^i , a_3 , b_2 couplings is essential for having large Majorana masses for right-handed neutrinos after symmetry breaking. Suppose that right-handed neutrinos ν^c , χ_{ν^c} , $\bar{\chi}_{\nu^c}$ acquire additional phases under transformations of the discrete symmetry:

$$\nu^c \rightarrow \nu^c e^{i\alpha}, \quad \chi_{\nu^c} \rightarrow \chi_{\nu^c} e^{i\beta}, \quad \bar{\chi}_{\nu^c} \rightarrow \bar{\chi}_{\nu^c} e^{i\gamma} \quad (19)$$

Then from the requirement that nonzero a_1^{ij} and b_1 are allowed while a_2^i , a_3 , b_2^i , c_2^i are set to zero by the symmetry:

$$\alpha + \gamma = \pi k, \quad \beta + \gamma = \frac{2\pi}{3}l, \quad (\alpha + \gamma) + (\beta + \gamma) \neq 2\pi m, \quad (20)$$

$$\beta + \gamma \neq \pi n, \quad (\alpha + \gamma) + 2(\beta + \gamma) \neq 2\pi q, \quad (\alpha + \gamma) + 3(\beta + \gamma) \neq 2\pi p.$$

Conditions (20) imply that $\frac{2}{3}l$ is not integer and, consequently, that $\alpha - \beta \neq 2\pi j$:

$$\alpha - \beta = (\alpha + \gamma) - (\beta + \gamma) = \pi(k - \frac{2}{3}l) \neq 2\pi j \neq 0 \quad (21)$$

In other words, ν^c and χ_{ν^c} have different transformation properties under the discrete symmetry. If the last term in (3) which is responsible for both small neutrino masses via the see-saw mechanism and leptogenesis is allowed by this symmetry, then the term $\chi_{\nu^c}(L_j H_k^u)$ is necessarily forbidden just as was assumed.


Bare mass term $M\chi_{\nu^c}\bar{\chi}_{\nu^c}$ is not invariant under transformations of the discrete symmetry and therefore is not allowed. From equations (20) it also follows that $\alpha + \gamma = \pi k$ with k — odd, so that bare mass term $M\nu^c\bar{\chi}_{\nu^c}$ is forbidden as well. Finally, coefficients b_3 and b_4 of the last two terms in (12) vanish for the same reasons.

After $\tilde{\chi}_{\nu^c}$ and $\tilde{\bar{\chi}}_{\nu^c}$ develop nonzero VEVs, the $U(1)$ symmetry, as well as the discrete symmetry (19), is broken down. Components of chiral (super)fields χ_{ν^c} , $\bar{\chi}_{\nu^c}$ and gauge (super)field become massive. As $m_{\chi}^2 = m_{\tilde{\chi}}^2$ is assumed, VEVs of $\tilde{\chi}_{\nu^c}$ and $\tilde{\bar{\chi}}_{\nu^c}$ are equal and it is natural to introduce new fields $h_1 = (\chi - \bar{\chi})/\sqrt{2}$ and $h_2 = (\chi + \bar{\chi})/\sqrt{2}$. The imaginary component of $\tilde{h}_1$ is ‘eaten up’ by the vector gauge field A , which acquires large mass $M_A = M = gqv_0 \sim 10^{14}$ GeV. The real component η of $\tilde{h}_1$ acquire the same mass $M_\eta = M$.

From the analysis of gauge interactions alone, it follows that two-component spinors λ (superpartner of A) and h_1 (superpartner of $\tilde{h}_1$) form a four-component Dirac spinor $M_f(h_1\lambda + \lambda^+h_1^+)$ with mass $M_f = M$. Non-renormalizable interactions induce Majorana-type mass term $\sim m_{soft}(h_1h_1 + c.c.)$. There is also Majorana-type mass term $\sim m_{soft}(\lambda\lambda + c.c.)$ coming from the soft supersymmetry breaking, so that two linear combinations of h_1 and λ are Majorana fermions with large ($\sim M$) and close masses. Non-renormalizable interactions (12) also induce masses $\sim m_{soft}$ for the real component of $\tilde{h}_2$ and its fermionic superpartner h_2 .

The gauge boson A , the fermion λ and the scalar η interact with other states in the fundamental representation of E_6 (in particular with Higgses) so that the self-energy of scalars receives large contributions from diagrams with exchange of these heavy fields. For instance, see the case of one loop in figure 1.

Figure 1. One-loop contributions to self-energy of scalars.



As has already been mentioned in the introduction, the supersymmetric structure of the Lagrangian ensures that all large corrections, associated with the exchange of heavy fields, cancel out and only terms proportional to the soft supersymmetry breaking parameters remain:

$$\Pi^0(p^2) = 4 g_{\omega_0}^2 q_{\tilde{\Phi}}^2 m_{\tilde{\Phi}}^2 B(m_{\tilde{\Phi}}^2, M^2), \quad B(x, y) = \frac{(2\pi\mu)^{2\epsilon}}{\pi^2} \int \frac{d^{4-2\epsilon}k}{[k^2 - x][(k - p)^2 - y]} \quad (22)$$

$SU_C(3) \times SU_L(2) \times SU_R(2) \times U^2(1)$ models. Tables 4 and 5 show, that it is sufficient to consider only the case of the first charge assignment and three $SU(3)_R \rightarrow SU_R(2)$ projections, the other cases being completely analogous.

In the case of the left-right symmetric (R) model, the zero modes in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ are $SU_R(2)$ doublets $\chi_R = (\chi_{e^c}, \chi_{\nu^c})$ and $\bar{\chi}_R = (\bar{\chi}_{\nu^c}, \bar{\chi}_{e^c})$. The scalar potential coming from gauge interactions (6) and soft supersymmetry breaking is of the form

$$V = \frac{1}{8} (g_{Y_L}^2 q_{Y_L}^2 + g_{Y_R}^2 q_{Y_R}^2) [(\tilde{\chi}_R^* \tilde{\chi}_R) - (\tilde{\bar{\chi}}_R^* \tilde{\bar{\chi}}_R)]^2 + \frac{g_{W_R}^2}{2} \left[\left(\tilde{\chi}_R^* \frac{\tau_i}{2} \tilde{\chi}_R \right) + \left(\tilde{\bar{\chi}}_R^* \frac{\tau_i}{2} \tilde{\bar{\chi}}_R \right) \right]^2 \quad (23)$$

$$- [m_{\chi}^2 (\tilde{\chi}_R^* \tilde{\chi}_R) + m_{\bar{\chi}}^2 (\tilde{\bar{\chi}}_R^* \tilde{\bar{\chi}}_R)]$$

Let $v_1 \equiv \langle \tilde{\chi}_{\nu^c} \rangle$, $v_2 \equiv \langle \tilde{\chi}_{e^c} \rangle$ and $u_1 \equiv \langle \tilde{\bar{\chi}}_{\nu^c} \rangle$, $u_2 \equiv \langle \tilde{\bar{\chi}}_{e^c} \rangle$. The corresponding classical potential is

$$V = \frac{1}{8} (g_{Y_L}^2 q_{Y_L}^2 + g_{Y_R}^2 q_{Y_R}^2) [(|v_1|^2 + |v_2|^2) - (|u_1|^2 + |u_2|^2)] \quad (24)$$

$$\begin{aligned}
 & + \frac{1}{2} g_{W_R}^2 |v_2^* v_1 + u_1^* u_2|^2 + \frac{1}{8} g_{W_R}^2 [(|v_2|^2 - |v_1|^2) + (|u_1|^2 - |u_2|^2)] \\
 & - [m_\chi^2 (|v_1|^2 + |v_2|^2) + m_{\bar{\chi}}^2 (|u_1|^2 + |u_2|^2)]
 \end{aligned}$$

There are symmetry breaking directions, which are $D - \text{flat}$. Vanishing of the first and the third term in (24) requires that $|v_1| = |u_1|$ and $|v_2| = |u_2|$. With these conditions satisfied, the second term in (24) vanishes if $v_1 = 0$ or $v_2 = 0$ or $\arg(u_2) + \arg(v_2) - \arg(u_1) - \arg(v_1) = \pi$.

The non-renormalizable superpotential, which bounds the classical potential from below, is similar to (12) with ν^c , χ_{ν^c} and $\bar{\chi}_{\nu^c}$ replaced with $SU_R(2)$ doublets $\psi_R = (e^c, \nu^c)$, $\chi_R = (\chi_{e^c}, \chi_{\nu^c})$ and $\bar{\chi}_R = (\bar{\chi}_{\nu^c}, \bar{\chi}_{e^c})$. Its explicit form is

$$\begin{aligned}
 W = M_c^{-1} [& a_1^{ij} (\psi_R^i \bar{\chi}_R) (\psi_R^j \bar{\chi}_R) + a_2^i (\psi_R^i \bar{\chi}_R) (\psi_R^i \bar{\chi}_R) + a_3 (\chi_R \bar{\chi}_R) (\chi_R \bar{\chi}_R)] \\
 & + M_c^{-3} [(\chi_R \bar{\chi}_R) (\chi_R \bar{\chi}_R) (\chi_R \bar{\chi}_R) + b^i (\psi_R^i \bar{\chi}_R) (\chi_R \bar{\chi}_R) (\chi_R \bar{\chi}_R) \\
 & + b_3^{ij} (\psi_R^i \bar{\chi}_R) (h^j \bar{\chi}_R) (\chi_R \bar{\chi}_R) + b_4^{ijk} (\psi_R^i \bar{\chi}_R) (\psi_R^j \bar{\chi}_R) (\psi_R^k \bar{\chi}_R)] + \dots
 \end{aligned} \tag{25}$$

The freedom of $SU_R(2)$ gauge transformations allows to rotate away a possible VEV for one of the isospin components of one of the scalar fields, so one can take $u_2 = 0$ at the minimum of the potential. Since the classical potential under consideration reaches its minimum in one of the $D - \text{flat}$ directions, v_2 is equal to zero as well. Then the following analysis is the same as in the case of the $SU_C(3) \times SU_L(2) \times U^3(1)$ model and furnishes the same result. The symmetry is broken down to $SU_C(3) \times SU_L(2) \times U^2(1)$.

The second skew left-right symmetric (R'') model differs from the one above in S instead of e^c being an $SU_{R''}(2)$ counterpart of the right-handed neutrino: $\chi_{R''} = (\chi_{\nu^c}, \chi_S)$ and $\bar{\chi}_{R''} = (\bar{\chi}_S, \bar{\chi}_{\nu^c})$. The symmetry is broken down to $SU_C(3) \times SU_L(2) \times U^2(1)$ as well.

In the case of the first skew left-right symmetric (R') model, the right-handed neutrino is an $SU_{R'}(2)$ singlet. Massless states in $\delta(27 + \overline{27})$ are χ_{ν^c} , $\chi_{R'} = (\chi_{H^u}, \chi_L)$ and $\bar{\chi}_{\nu^c}$, $\bar{\chi}_{R'} = (\bar{\chi}_L, \bar{\chi}_{H^u})$. The classical potential coming from renormalizable and nonrenormalizable interactions is similar to (15) and yields the same results. The symmetry is broken down to $SU_C(3) \times SU_L(2) \times SU_{R'}(2) \times U(1)$.

Since the Higgs doublets (χ_{H^u}, χ_L) and $(\bar{\chi}_L, \bar{\chi}_{H^u})$ are contained in $\delta(\mathbf{27} + \overline{\mathbf{27}})$ as zero modes, there are directions in which the D -term potential vanishes for whatever large VEVs of these fields, i.e., there is a risk of breaking electroweak symmetry at a very high scale. To avoid it, the part of the classical potential which comes from the soft supersymmetry breaking should be positive in those directions, as is the case in the MSSM.

The coexistence of all the terms in the second row of (3) leads to the rapid proton decay, mediated by new D and D^c quarks, unless those are very heavy. VEV of S which gives masses $\lambda_5^{ijk} \langle S_i \rangle$ to D and D^c is also the source of μ -term $\lambda_4^{ijk} \langle S_i \rangle$. Although neither for $G_{U(1)}$ nor for $G_{SU_R(2)}$ the couplings λ_5^{ijk} and λ_4^{ijk} are related by symmetry, it is not natural to expect D and D^c to be much heavier than 1 TeV which is insufficient to

suppress the proton decay. A solution to this problem may be provided by an appropriate discrete symmetry, which forbids some of the couplings in (3).

Table 6. Transformation properties of components of $SU_R(2)$ doublets.

Model	R				R'								R''					
	d^c	u^c	H^u	H^d	e^c	ν^c	D^c	u^c	H^u	L	e^c	S	D^c	d^c	H^d	L	ν^c	S
1	+	+	+	+	-	-	-	+	+	-	-	+	-	+	+	-	-	+
2	+	+	+	+	-	-	+	+	+	-	-	+	+	+	+	-	-	+

If this is a Z_2 symmetry, then there are only two models [17, 20] compatible with leptogenesis and nonzero neutrino masses. In the first model L and e^c , ν^c , D , D^c are odd, while the rest of the states are even, so that $\lambda_9 = \lambda_{10} = 0$. The second model differs from the first in D and D^c being even, so that $\lambda_6 = \lambda_7 = \lambda_8 = 0$. Transformations of the discrete symmetry should commute with transformations of the gauge symmetry. Table 6 shows, that this condition is satisfied only in the case of the left–right symmetric (R) model, while in both skew left–right symmetric models components of $SU_R(2)$ doublets transform differently. Therefore, if proton stability is assured by such a discrete symmetry, the only allowed gauge group after breaking of G is $G'_{U(1)} = SU_C(3) \times SU_L(2) \times U^2(1)$.

5. Conclusions

In this paper, an attempt to derive ‘low–energy’ extensions of the SM, compatible with the Fukugita–Yanagida scenario, from a superstring inspired E_6 model has been made.

This model allows for six charge assignments compatible with the standard model. Charge conservation in the processes involving states of different generations requires that the same charge assignment be used for all generations.

As is discussed above, the gauge symmetry is broken in a sequence of stages. The first stage is due to Calabi–Yau compactification and the effective Higgs mechanism. The condition that the proton is long-lived requires that the symmetry is broken either to $G_{U(1)}$ or $G_{SU_R(2)}$.

As the temperature goes down, supersymmetry breaks down spontaneously in the hidden sector. Supersymmetry breaking is mediated to the visible sector by gravity and manifests itself in the soft terms.

At the next stage right–handed scalar neutrinos and right–handed neutrinos of two additional generations ($\mathbf{27} + \overline{\mathbf{27}}$) develop nonzero VEV, breaking the $B - L$ symmetry. The introduced simple discrete symmetry ensures that $B - L$ is broken at the scale which is sufficiently high for generating large masses of the right–handed neutrinos and that right–handed scalar neutrinos of the three known generations do not acquire VEV. It also forbids Yukawa couplings which, if present, would induce large masses of the conventional neutrinos. The supersymmetric structure of the theory ensures that large

quantum corrections to masses of scalars, associated with the presence of heavy gauge fields, cancel out. Provided that a rapid proton decay mediated by the new quarks is forbidden by a Z_2 symmetry, the sought ‘low-energy’ gauge group is given by $G'_{U(1)}$ for both allowed intermediate scale gauge groups $G_{U(1)}$ and $G_{SU_R(2)}$.

Apart from the additional (compared to the SM) $U(1)$ symmetry, the characteristic feature of this ‘low-energy’ model is the extended particle content. In addition to the known particles and right-handed neutrinos it contains the SM singlets S , new heavy quarks and three generations of Higgses, as well as their superpartners. Since both new quarks and Higgses couple to right-handed neutrinos, there are more $B - L$ violating decay channels than in the SM or its supersymmetric extension. At the same time there are more processes which wash the generated $B - L$ asymmetry out. The interplay of these effects requires further investigation.

Acknowledgments

I would like to thank Professor E A Paschos for inspiration and useful discussions. Financial support from the Graduiertenkolleg 841 Physik der Elementarteilchen an Beschleunigern und im Universum at Dortmund University is gratefully acknowledged.

References

- [1] A.D.Sakharov, Pis'ma Zh.Eksp. Teor. Fiz. **5**, 32 (1967).
- [2] G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976).
- [3] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. **B155**, 36 (1985).
- [4] M. Fukugita and T. Yanagida, Phys. Lett. **B174**, 45 (1986).
- [5] J. Orloff, hep-ph/0307351.
- [6] A. S. Joshipura and E. A. Paschos, hep-ph/9906498.
- [7] A. S. Joshipura, E. A. Paschos and W. Rodejohann, JHEP **08**, 029 (2001), [hep-ph/0105175].
- [8] A. S. Joshipura, E. A. Paschos and W. Rodejohann, Nucl. Phys. **B611**, 227 (2001), [hep-ph/0104228].
- [9] T. Matsuoka and D. Suematsu, Prog. Theor. Phys. **76**, 886 (1986).
- [10] E. Witten, Nucl. Phys. **B258**, 75 (1985).
- [11] E. Dynkin, Am. Math. Soc. Trans., Ser. 2 **6**, 245 (1957).
- [12] R. Slansky, Phys. Rept. **79**, 1 (1981).
- [13] M. Plumacher, hep-ph/9807557.
- [14] G. W. Anderson and T. Blazek, J. Math. Phys. **41**, 4808 (2000), [hep-ph/9912365].
- [15] Y. Hosotani, Phys. Lett. **B126**, 309 (1983).
- [16] T. Matsuoka and D. Suematsu, Nucl. Phys. **B274**, 106 (1986).
- [17] T. Hambye, E. Ma, M. Raidal and U. Sarkar, Phys. Lett. **B512**, 373 (2001), [hep-ph/0011197].
- [18] M. Dine, V. Kaplunovsky, M. L. Mangano, C. Nappi and N. Seiberg, Nucl. Phys. **B259**, 549 (1985).
- [19] C. A. Lutken and G. G. Ross, Phys. Lett. **B214**, 357 (1988).
- [20] E. Ma, Phys. Rev. Lett. **60**, 1363 (1988).

Table 7. Weights of $\mathbf{27}$ of E_6 . Here $\varepsilon = \frac{1}{3}(\alpha + \beta + \gamma)$, $\bar{x} \equiv -x$. The zero root breaking SM gauge group preserving direction is given by $Z = (-\varepsilon, \varepsilon, \beta, 2\varepsilon - \gamma, \varepsilon, 0)$.

E_6	$SU(3)$			$SU(2)_Y$				Assignments						(Z, ρ)
	C	L	R	L	R	R'	R''	I	II	III	IV	V	VI	
27	3	3	1	2										
[100000]	[10]	[10]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	u	u	u	u	u	u	$-\varepsilon$
[110010]	[11]	[10]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	u	u	u	u	u	u	$-\varepsilon$
[100001]	[01]	[10]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	u	u	u	u	u	u	$-\varepsilon$
[000011]	[10]	[11]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	d	d	d	d	d	d	$-\varepsilon$
[010001]	[11]	[11]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	d	d	d	d	d	d	$-\varepsilon$
[000010]	[01]	[11]	[00]	[1] _{,1}	[0] _{,0}	[0] _{,0}	[0] _{,0}	d	d	d	d	d	d	$-\varepsilon$
27	3	3	1	1										
[110000]	[10]	[01]	[00]	[0] _{,2}	[0] _{,0}	[0] _{,0}	[0] _{,0}	D	D	D	D	D	D	2ε
[100010]	[11]	[01]	[00]	[0] _{,2}	[0] _{,0}	[0] _{,0}	[0] _{,0}	D	D	D	D	D	D	2ε
[110001]	[01]	[01]	[00]	[0] _{,2}	[0] _{,0}	[0] _{,0}	[0] _{,0}	D	D	D	D	D	D	2ε
27	3	1	3	1										
[001101]	[01]	[00]	[10]	[0] _{,0}	[1] _{,1}	[1] _{,1}	[0] _{,2}	u^c	u^c	d^c	D^c	d^c	D^c	$-\varepsilon + \alpha$
[011100]	[11]	[00]	[10]	[0] _{,0}	[1] _{,1}	[1] _{,1}	[0] _{,2}	u^c	u^c	d^c	D^c	d^c	D^c	$-\varepsilon + \alpha$
[001100]	[10]	[00]	[10]	[0] _{,0}	[1] _{,1}	[1] _{,1}	[0] _{,2}	u^c	u^c	d^c	D^c	d^c	D^c	$-\varepsilon + \alpha$
[000111]	[01]	[00]	[01]	[0] _{,0}	[0] _{,2}	[1] _{,1}	[1] _{,1}	D^c	d^c	D^c	d^c	u^c	u^c	$-\varepsilon + \gamma$
[010100]	[11]	[00]	[01]	[0] _{,0}	[0] _{,2}	[1] _{,1}	[1] _{,1}	D^c	d^c	D^c	d^c	u^c	u^c	$-\varepsilon + \gamma$
[000110]	[10]	[00]	[01]	[0] _{,0}	[0] _{,2}	[1] _{,1}	[1] _{,1}	D^c	d^c	D^c	d^c	u^c	u^c	$-\varepsilon + \gamma$
[011000]	[01]	[00]	[11]	[0] _{,0}	[1] _{,1}	[0] _{,2}	[1] _{,1}	d^c	D^c	u^c	u^c	D^c	d^c	$-\varepsilon + \beta$
[001011]	[11]	[00]	[11]	[0] _{,0}	[1] _{,1}	[0] _{,2}	[1] _{,1}	d^c	D^c	u^c	u^c	D^c	d^c	$-\varepsilon + \beta$
[011001]	[10]	[00]	[11]	[0] _{,0}	[1] _{,1}	[0] _{,2}	[1] _{,1}	d^c	D^c	u^c	u^c	D^c	d^c	$-\varepsilon + \beta$
27	1	3	3	2										
[001111]	[00]	[11]	[10]	[1] _{,1}	[1] _{,1}	[1] _{,1}	[0] _{,2}	E^c	E^c	N	ν	N	ν	$2\varepsilon - \alpha$
[101100]	[00]	[10]	[10]	[1] _{,1}	[1] _{,1}	[1] _{,1}	[0] _{,2}	N^c	N^c	E	e	E	e	$2\varepsilon - \alpha$
[011010]	[00]	[11]	[11]	[1] _{,1}	[1] _{,1}	[0] _{,2}	[1] _{,1}	N	ν	E^c	E^c	ν	N	$2\varepsilon - \beta$
[111001]	[00]	[10]	[11]	[1] _{,1}	[1] _{,1}	[0] _{,2}	[1] _{,1}	E	e	N^c	N^c	e	E	$2\varepsilon - \beta$
[000101]	[00]	[11]	[01]	[1] _{,1}	[0] _{,2}	[1] _{,1}	[1] _{,1}	ν	N	ν	N	E^c	E^c	$2\varepsilon - \gamma$
[100110]	[00]	[10]	[01]	[1] _{,1}	[0] _{,2}	[1] _{,1}	[1] _{,1}	e	E	e	E	N^c	N^c	$2\varepsilon - \gamma$
27	1	3	3	1										
[111100]	[00]	[01]	[10]	[0] _{,2}	[1] _{,1}	[1] _{,1}	[0] _{,2}	e^c	e^c	ν^c	S	ν^c	S	$-\varepsilon - \alpha$
[101001]	[00]	[01]	[11]	[0] _{,2}	[1] _{,1}	[0] _{,2}	[1] _{,1}	ν^c	S	e^c	e^c	S	ν^c	$-\varepsilon - \beta$
[110110]	[00]	[01]	[01]	[0] _{,2}	[0] _{,2}	[1] _{,1}	[1] _{,1}	S	ν^c	S	ν^c	e^c	e^c	$-\varepsilon - \gamma$

Table 8. Weights of **78** of E_6 (beginning). Color, isospin and hypercharge (as well as $B - L$, unless otherwise indicated) of the states designated like the SM states are the same as charges of their SM counterparts. If present, subscript is trice the $B - L$ charge. All charges of overlined states are opposite to those of non overlined. For example, for $\overline{d}_2$: $I_3 = \frac{1}{2}$, $Y = -\frac{1}{3}$, $B - L = \frac{2}{3}$.

E_6	$SU(3)$			$SU(2)_Y$				Assignments						(Z, ρ)
	C	L	R	L	R	R'	R''	I	II	III	IV	V	VI	
78	1	8	1											
$[2\overline{1}0000]$	$[00]$	$[11]$	$[00]$	$[1]_{,3}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$\overline{e}$	$\overline{e}$	$\overline{e}$	$\overline{e}$	$\overline{e}$	$\overline{e}$	-3ε
$[1\overline{1}00\overline{1}1]$	$[00]$	$[\overline{1}2]$	$[00]$	$[\overline{1}]_{,3}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$\overline{\nu}$	$\overline{\nu}$	$\overline{\nu}$	$\overline{\nu}$	$\overline{\nu}$	$\overline{\nu}$	-3ε
$[10001\overline{1}]$	$[00]$	$[2\overline{1}]$	$[00]$	$[2]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	W^+	W^+	W^+	W^+	W^+	W^+	0
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	W^0	W^0	W^0	W^0	W^0	W^0	0
$[\overline{1}000\overline{1}1]$	$[00]$	$[\overline{2}1]$	$[00]$	$[\overline{2}]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	W^-	W^-	W^-	W^-	W^-	W^-	0
$[\overline{1}1001\overline{1}]$	$[00]$	$[1\overline{2}]$	$[00]$	$[1]_{,\overline{3}}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	ν	ν	ν	ν	ν	ν	3ε
$[210000]$	$[00]$	$[\overline{1}\overline{1}]$	$[00]$	$[\overline{1}]_{,3}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	e	e	e	e	e	e	3ε
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	γ^0	γ^0	γ^0	γ^0	γ^0	γ^0	0
78	1	1	8											
$[001\overline{2}10]$	$[00]$	$[00]$	$[11]$	$[0]_{,0}$	$[1]_{,3}$	$[2]_{,0}$	$[1]_{,\overline{3}}$	e^c	$\overline{e_0^c}$	ν^c	$\overline{\nu^c}$	$\overline{e_0^c}$	$\overline{e^c}$	$\gamma - \alpha$
$[01\overline{1}\overline{1}11]$	$[00]$	$[00]$	$[\overline{1}2]$	$[0]_{,0}$	$[\overline{1}]_{,3}$	$[1]_{,\overline{3}}$	$[2]_{,0}$	ν^c	$\overline{\nu^c}$	e^c	$\overline{e_0^c}$	$\overline{e^c}$	$\overline{e_0^c}$	$\gamma - \beta$
$[0\overline{1}2\overline{1}0\overline{1}]$	$[00]$	$[00]$	$[2\overline{1}]$	$[0]_{,0}$	$[2]_{,0}$	$[1]_{,3}$	$[\overline{1}]_{,\overline{3}}$	$\overline{e_0^c}$	$\overline{e^c}$	$\overline{e_0^c}$	$\overline{e^c}$	ν^c	$\overline{\nu^c}$	$\beta - \alpha$
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	ϕ^0	ϕ^0	ϕ^0	ϕ^0	ϕ^0	ϕ^0	0
$[01\overline{2}101]$	$[00]$	$[00]$	$[2\overline{1}]$	$[0]_{,0}$	$[2]_{,0}$	$[\overline{1}]_{,\overline{3}}$	$[1]_{,3}$	$\overline{e_0^c}$	$\overline{e^c}$	e_0^c	e^c	$\overline{\nu^c}$	ν^c	$\alpha - \beta$
$[0\overline{1}11\overline{1}\overline{1}]$	$[00]$	$[00]$	$[12]$	$[0]_{,0}$	$[1]_{,\overline{3}}$	$[\overline{1}]_{,3}$	$[2]_{,0}$	$\overline{\nu^c}$	ν^c	$\overline{e^c}$	$\overline{e_0^c}$	e^c	e_0^c	$\beta - \gamma$
$[00\overline{1}2\overline{1}0]$	$[00]$	$[00]$	$[\overline{1}\overline{1}]$	$[0]_{,0}$	$[\overline{1}]_{,3}$	$[2]_{,0}$	$[\overline{1}]_{,3}$	$\overline{e^c}$	$\overline{e_0^c}$	$\overline{\nu^c}$	ν^c	e_0^c	e^c	$\alpha - \gamma$
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	ω^0	ω^0	ω^0	ω^0	ω^0	ω^0	0
78	8	1	1											
$[000001]$	$[11]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[0\overline{1}0011]$	$[\overline{1}2]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[0100\overline{1}0]$	$[2\overline{1}]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[000000]$	$[00]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[0100\overline{1}\overline{1}]$	$[1\overline{2}]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[0\overline{1}0010]$	$[\overline{2}1]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
$[00000\overline{1}]$	$[\overline{1}\overline{1}]$	$[00]$	$[00]$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	$[0]_{,0}$	g	g	g	g	g	g	0
78	$\overline{3}$	3	3											
$[1\overline{1}1\overline{1}10]$	$[01]$	$[10]$	$[10]$	$[1]_{,1}$	$[1]_{,1}$	$[1]_{,1}$	$[0]_{,2}$	$\overline{X}$	$\overline{X}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$-\alpha$
$[101\overline{1}0\overline{1}]$	$[\overline{1}\overline{1}]$	$[10]$	$[10]$	$[1]_{,1}$	$[1]_{,1}$	$[1]_{,1}$	$[0]_{,2}$	$\overline{X}$	$\overline{X}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$-\alpha$
$[1\overline{1}1\overline{1}1\overline{1}]$	$[\overline{1}0]$	$[10]$	$[10]$	$[1]_{,1}$	$[1]_{,1}$	$[1]_{,1}$	$[0]_{,2}$	$\overline{X}$	$\overline{X}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$-\alpha$
$[10\overline{1}011]$	$[01]$	$[10]$	$[\overline{1}\overline{1}]$	$[1]_{,1}$	$[\overline{1}]_{,1}$	$[0]_{,2}$	$[1]_{,1}$	$\overline{d_2}$	$\overline{d}$	$\overline{X}$	$\overline{X}$	$\overline{d}$	$\overline{d_2}$	$-\beta$
$[11\overline{1}000]$	$[\overline{1}\overline{1}]$	$[10]$	$[\overline{1}\overline{1}]$	$[1]_{,1}$	$[\overline{1}]_{,1}$	$[0]_{,2}$	$[1]_{,1}$	$\overline{d_2}$	$\overline{d}$	$\overline{X}$	$\overline{X}$	$\overline{d}$	$\overline{d_2}$	$-\beta$
$[10\overline{1}010]$	$[\overline{1}0]$	$[10]$	$[\overline{1}\overline{1}]$	$[1]_{,1}$	$[\overline{1}]_{,1}$	$[0]_{,2}$	$[1]_{,1}$	$\overline{d_2}$	$\overline{d}$	$\overline{X}$	$\overline{X}$	$\overline{d}$	$\overline{d_2}$	$-\beta$
$[1\overline{1}0100]$	$[01]$	$[10]$	$[0\overline{1}\overline{1}]$	$[1]_{,1}$	$[0]_{,2}$	$[\overline{1}]_{,1}$	$[\overline{1}]_{,1}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{X}$	$\overline{X}$	$-\gamma$
$[1001\overline{1}\overline{1}]$	$[\overline{1}\overline{1}]$	$[10]$	$[0\overline{1}\overline{1}]$	$[1]_{,1}$	$[0]_{,2}$	$[\overline{1}]_{,1}$	$[\overline{1}]_{,1}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{X}$	$\overline{X}$	$-\gamma$
$[1\overline{1}010\overline{1}]$	$[\overline{1}0]$	$[10]$	$[0\overline{1}\overline{1}]$	$[1]_{,1}$	$[0]_{,2}$	$[\overline{1}]_{,1}$	$[\overline{1}]_{,1}$	$\overline{d}$	$\overline{d_2}$	$\overline{d}$	$\overline{d_2}$	$\overline{X}$	$\overline{X}$	$-\gamma$
$[0\overline{1}1\overline{1}0\overline{1}]$	$[01]$	$[\overline{1}\overline{1}]$	$[10]$	$[\overline{1}]_{,1}$	$[1]_{,1}$	$[1]_{,1}$	$[0]_{,2}$	$\overline{Y}$	$\overline{Y}$	$\overline{u_2}$	$\overline{u}$	$\overline{u_2}$	$\overline{u}$	$-\alpha$

Table 8. Weights of **78** of E_6 (continuation).

78	$\bar{3}$	3	3											
[001 $\bar{1}\bar{1}$ 0]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[10]	[$\bar{1}$] _{,1}	[1] _{,1}	[1] _{,1}	[0] _{,2}	$\bar{Y}$	$\bar{Y}$	$\bar{u}_2$	$\bar{u}$	$\bar{u}_2$	$\bar{u}$	$-\alpha$
[0 $\bar{1}$ 1 $\bar{1}$ 00]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[10]	[$\bar{1}$] _{,1}	[1] _{,1}	[1] _{,1}	[0] _{,2}	$\bar{Y}$	$\bar{Y}$	$\bar{u}_2$	$\bar{u}$	$\bar{u}_2$	$\bar{u}$	$-\alpha$
[00 $\bar{1}$ 002]	[01]	[$\bar{1}$ 1]	[$\bar{1}$ 1]	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	$\bar{u}_2$	$\bar{u}$	$\bar{Y}$	$\bar{Y}$	$\bar{u}$	$\bar{u}_2$	$-\beta$
[01 $\bar{1}$ 0 $\bar{1}$ 1]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[$\bar{1}$ 1]	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	$\bar{u}_2$	$\bar{u}$	$\bar{Y}$	$\bar{Y}$	$\bar{u}$	$\bar{u}_2$	$-\beta$
[00 $\bar{1}$ 001]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[$\bar{1}$ 1]	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	$\bar{u}_2$	$\bar{u}$	$\bar{Y}$	$\bar{Y}$	$\bar{u}$	$\bar{u}_2$	$-\beta$
[0 $\bar{1}$ 01 $\bar{1}$ 1]	[01]	[$\bar{1}$ 1]	[0 $\bar{1}$]	[$\bar{1}$] _{,1}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	$\bar{u}$	$\bar{u}_2$	$\bar{u}$	$\bar{u}_2$	$\bar{Y}$	$\bar{Y}$	$-\gamma$
[0001 $\bar{2}$ 0]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[0 $\bar{1}$]	[$\bar{1}$] _{,1}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	$\bar{u}$	$\bar{u}_2$	$\bar{u}$	$\bar{u}_2$	$\bar{Y}$	$\bar{Y}$	$-\gamma$
[0 $\bar{1}$ 01 $\bar{1}$ 0]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[0 $\bar{1}$]	[$\bar{1}$] _{,1}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	$\bar{u}$	$\bar{u}_2$	$\bar{u}$	$\bar{u}_2$	$\bar{Y}$	$\bar{Y}$	$-\gamma$
[1 $\bar{1}$ 0 $\bar{1}$ 10]	[01]	[0 $\bar{1}$]	[10]	[0] _{,2}	[1] _{,1}	[1] _{,1}	[0] _{,2}	d^c	d^c	u^c	u_4^c	u^c	u_4^c	$\beta + \gamma$
[$\bar{1}$ 1 $\bar{1}$ 0 $\bar{1}$]	[1 $\bar{1}$]	[0 $\bar{1}$]	[10]	[0] _{,2}	[1] _{,1}	[1] _{,1}	[0] _{,2}	d^c	d^c	u^c	u_4^c	u^c	u_4^c	$\beta + \gamma$
[1 $\bar{1}$ 0 $\bar{1}$ 1 $\bar{1}$]	[$\bar{1}$ 0]	[0 $\bar{1}$]	[10]	[0] _{,2}	[1] _{,1}	[1] _{,1}	[0] _{,2}	d^c	d^c	u^c	u_4^c	u^c	u_4^c	$\beta + \gamma$
[$\bar{1}$ 1 $\bar{1}$ 011]	[01]	[0 $\bar{1}$]	[$\bar{1}$ 1]	[0] _{,2}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	u^c	u_4^c	d^c	d^c	u_4^c	u^c	$\alpha + \gamma$
[1 $\bar{2}$ 1000]	[1 $\bar{1}$]	[0 $\bar{1}$]	[$\bar{1}$ 1]	[0] _{,2}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	u^c	u_4^c	d^c	d^c	u_4^c	u^c	$\alpha + \gamma$
[$\bar{1}$ 1 $\bar{1}$ 010]	[$\bar{1}$ 0]	[0 $\bar{1}$]	[$\bar{1}$ 1]	[0] _{,2}	[$\bar{1}$] _{,1}	[0] _{,2}	[1] _{,1}	u^c	u_4^c	d^c	d^c	u_4^c	u^c	$\alpha + \gamma$
[$\bar{1}$ 00100]	[01]	[0 $\bar{1}$]	[0 $\bar{1}$]	[0] _{,2}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	u_4^c	u^c	u_4^c	u^c	d^c	d^c	$\alpha + \beta$
[$\bar{1}$ 101 $\bar{1}$ 1]	[1 $\bar{1}$]	[0 $\bar{1}$]	[0 $\bar{1}$]	[0] _{,2}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	u_4^c	u^c	u_4^c	u^c	d^c	d^c	$\alpha + \beta$
[$\bar{1}$ 0010 $\bar{1}$]	[$\bar{1}$ 0]	[0 $\bar{1}$]	[0 $\bar{1}$]	[0] _{,2}	[0] _{,2}	[$\bar{1}$] _{,1}	[$\bar{1}$] _{,1}	u_4^c	u^c	u_4^c	u^c	d^c	d^c	$\alpha + \beta$
78	3	$\bar{3}$	$\bar{3}$											
[100 $\bar{1}$ 01]	[10]	[01]	[01]	[0] _{,2}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{d}^c$	$\bar{d}^c$	$-\alpha - \beta$
[1 $\bar{1}$ 0 $\bar{1}$ 11]	[$\bar{1}$ 1]	[01]	[01]	[0] _{,2}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{d}^c$	$\bar{d}^c$	$-\alpha - \beta$
[100 $\bar{1}$ 00]	[0 $\bar{1}$]	[01]	[01]	[0] _{,2}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{d}^c$	$\bar{d}^c$	$-\alpha - \beta$
[1 $\bar{1}$ 10 $\bar{1}$ 0]	[10]	[01]	[$\bar{1}$ 1]	[0] _{,2}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$-\alpha - \gamma$
[1 $\bar{2}$ 1000]	[$\bar{1}$ 1]	[01]	[$\bar{1}$ 1]	[0] _{,2}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$-\alpha - \gamma$
[1 $\bar{1}$ 10 $\bar{1}$ 1]	[0 $\bar{1}$]	[01]	[$\bar{1}$ 1]	[0] _{,2}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$-\alpha - \gamma$
[10 $\bar{1}$ 1 $\bar{1}$ 1]	[10]	[01]	[$\bar{1}$ 0]	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$-\beta - \gamma$
[1 $\bar{1}$ 1101]	[$\bar{1}$ 1]	[01]	[$\bar{1}$ 0]	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$-\beta - \gamma$
[10 $\bar{1}$ 1 $\bar{1}$ 0]	[0 $\bar{1}$]	[01]	[$\bar{1}$ 0]	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	$\bar{d}^c$	$\bar{d}^c$	$\bar{u}^c$	$\bar{u}_4^c$	$\bar{u}^c$	$\bar{u}_4^c$	$-\beta - \gamma$
[010 $\bar{1}$ 10]	[10]	[1 $\bar{1}$]	[01]	[1] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	u	u_2	u	u_2	Y	Y	γ
[000 $\bar{1}$ 20]	[$\bar{1}$ 1]	[1 $\bar{1}$]	[01]	[1] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	u	u_2	u	u_2	Y	Y	γ
[010 $\bar{1}$ 1 $\bar{1}$]	[0 $\bar{1}$]	[1 $\bar{1}$]	[01]	[1] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	u	u_2	u	u_2	Y	Y	γ
[00100 $\bar{1}$]	[10]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	u_2	u	Y	Y	u	u_2	β
[0 $\bar{1}$ 101 $\bar{1}$]	[$\bar{1}$ 1]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	u_2	u	Y	Y	u	u_2	β
[00100 $\bar{2}$]	[0 $\bar{1}$]	[1 $\bar{1}$]	[$\bar{1}$ 1]	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	u_2	u	Y	Y	u	u_2	β
[01 $\bar{1}$ 100]	[10]	[1 $\bar{1}$]	[$\bar{1}$ 0]	[1] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	Y	Y	u_2	u	u_2	u	α
[00 $\bar{1}$ 110]	[$\bar{1}$ 1]	[1 $\bar{1}$]	[$\bar{1}$ 0]	[1] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	Y	Y	u_2	u	u_2	u	α
[01 $\bar{1}$ 10 $\bar{1}$]	[0 $\bar{1}$]	[1 $\bar{1}$]	[$\bar{1}$ 0]	[1] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	Y	Y	u_2	u	u_2	u	α
[$\bar{1}$ 10 $\bar{1}$ 01]	[10]	[$\bar{1}$ 0]	[01]	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	d	d_2	d	d_2	X	X	γ
[$\bar{1}$ 00 $\bar{1}$ 11]	[$\bar{1}$ 1]	[$\bar{1}$ 0]	[01]	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	d	d_2	d	d_2	X	X	γ
[$\bar{1}$ 10 $\bar{1}$ 00]	[0 $\bar{1}$]	[$\bar{1}$ 0]	[01]	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	[1] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	d	d_2	d	d_2	X	X	γ
[$\bar{1}$ 010 $\bar{1}$ 0]	[10]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[$\bar{1}$] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	d_2	d	X	X	d	d_2	β
[$\bar{1}$ 11000]	[$\bar{1}$ 1]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[$\bar{1}$] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	d_2	d	X	X	d	d_2	β
[$\bar{1}$ 010 $\bar{1}$ 1]	[0 $\bar{1}$]	[$\bar{1}$ 0]	[$\bar{1}$ 1]	[$\bar{1}$] _{,$\bar{1}$}	[1] _{,$\bar{1}$}	[0] _{,2}	[$\bar{1}$] _{,$\bar{1}$}	d_2	d	X	X	d	d_2	β
[$\bar{1}$ 1 $\bar{1}$ 1 $\bar{1}$ 1]	[10]	[$\bar{1}$ 0]	[$\bar{1}$ 0]	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	X	X	d_2	d	d_2	d	α
[$\bar{1}$ 0 $\bar{1}$ 101]	[$\bar{1}$ 1]	[$\bar{1}$ 0]	[$\bar{1}$ 0]	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	X	X	d_2	d	d_2	d	α
[$\bar{1}$ 1 $\bar{1}$ 1 $\bar{1}$ 0]	[0 $\bar{1}$]	[$\bar{1}$ 0]	[$\bar{1}$ 0]	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[$\bar{1}$] _{,$\bar{1}$}	[0] _{,2}	X	X	d_2	d	d_2	d	α